

$\gamma^*N\Delta$ Form Factors from a Relativistic Dynamical Model of Pion Electroproduction

G. L. Caia,^{1,*} V. Pascalutsa,^{2,3,†} J. A. Tjon,^{3,4,‡} and L. E. Wright^{1,§}

¹ *Institute of Nuclear and Particle Physics (INPP),
Department of Physics and Astronomy,
Ohio University, Athens, OH 45701*

² *Department of Physics, The College of William & Mary, Williamsburg, VA 23188*

³ *Theory Group, Jefferson Laboratory,
12000 Jefferson Ave, Newport News, VA 23606*

⁴ *Department of Physics, University of Maryland, College Park, MD 20742*

Abstract

We obtain the electromagnetic form factors of the $\gamma N\Delta$ transition by analyzing recent pion-electroproduction data using a fully relativistic dynamical model. Special care is taken to satisfy Ward-Takahashi identities for the Born term in the presence of form factors thereby allowing the use of realistic electromagnetic form factors of the nucleon and pion. We parametrize the Q^2 dependence of the *bare* $\gamma N\Delta$ form factors by a three-parameter form which is consistent with the asymptotic behavior inferred from QCD. The parameters of the bare $\gamma N\Delta$ form factors are the only free parameters of the model and are fitted to the differential cross-section and multipole-analysis data up to $Q^2 = 4$ (GeV/c)² in the $\Delta(1232)$ -resonance region. This analysis emphasizes the significance of the pion-cloud effects in the extraction of the resonance parameters.

PACS numbers: 12.38.Aw, 13.40.Gp, 13.60.Le, 14.20.Gk

*Electronic address: caia@phy.ohiou.edu

†Electronic address: vlad@jlab.org

‡Electronic address: tjon@jlab.org

§Electronic address: wright@ohiou.edu

Recently a covariant quasipotential description of pion-nucleon (πN) interaction [1] has been extended to the process of pion photo-production [2]. Here we further extended this description to the case of pion *electro*-production and use the resulting model to extract the electromagnetic N to Δ transition form factors. In doing so we exploit the results of the recent pion production experiments at LEGS [3], MAMI [4], JLab [5, 6], and MIT-Bates [7], as well as the multipole analysis MAID [8].

One of the aims of our investigation is to disentangle the effect of the resonance excitation from the competing background mechanisms, such as pion rescattering in the final state. Admittedly, any such separation of resonance and background is model dependent. We have attempted to constrain this model-dependence by adhering to general principles, such as relativistic covariance, current conservation, unitarity, chiral symmetry. The model is based on a πN - γN coupled-channel equation which when solved to the first order in the electromagnetic coupling e leads to the electroproduction amplitude, $T_{\pi\gamma^*} = V_{\pi\gamma^*} + T_{\pi\pi} G_\pi V_{\pi\gamma^*}$, where $V_{\pi\gamma^*}$ is a basic electroproduction potential, G_π is the pion-nucleon propagator and $T_{\pi\pi}$ is the full πN amplitude. Thus, the pion rescattering effects are included as the final state interaction. The πN amplitude satisfies an integral equation on its own. The details on reconstructing this amplitude and its fit to the πN elastic scattering are presented in [1].

Our model potential for the pion electro-production is shown in Fig. 1. It includes the Born term (we use the pseudo-vector πNN coupling required by chiral symmetry considerations), the t -channel exchange of ρ - and ω - mesons, and the Δ -isobar exchange.

The πN final state interaction dresses the s -channel nucleon and resonance contributions, leading in particular to the mass, field and coupling constant renormalizations. Therefore, both N - and Δ -pole contributions in $V_{\pi\gamma}$ are included using the *bare* mass and coupling parameters obtained from the equation for the πN amplitude. The renormalization conditions together with unitarity demand that the same propagators and πN vertices, including the cutoff functions, appear in both the πN and γN potentials. Thus, all these ingredients are fixed by the analysis of πN scattering [1].

On the other hand, the electromagnetic interaction is constrained by the electromagnetic *gauge invariance*. At this point one is often concerned with the problem of how to introduce the electromagnetic form factors for nucleon and pion in a way consistent with gauge invariance. A common, but not viable, solution to this problem, implemented for instance in Refs. [8, 9, 10], is to choose all of the electromagnetic form factors that go into the Born term (i.e., nucleon, pion and axial form factors) to be *the same*. This prescription does enforce the current conservation, however the Ward-Takahashi (WT) identities cannot be satisfied in this way. Furthermore, it is clear that the requirement of gauge invariance should not be able to restrict the Q^2 behavior of the electromagnetic interaction (see, *e.g.*, Ref. [11]). Finally, it is well known, especially in view of the new JLab experiments [13], that these form factors *are not* the same.

For these reasons we sought for a solution that permits arbitrary choices of the electromagnetic form factors, yet is fully consistent with gauge invariance. Following the arguments given in, *e.g.*, [11, 12], we find that an arbitrary form factor $F(Q^2)$ can be accommodated by the following replacement the current:

$$J^\mu \rightarrow J'^\mu(Q^2) = J^\mu + [F(Q^2) - 1] O^{\mu\nu} J_\nu, \quad (1)$$

where $O^{\mu\nu} = g^{\mu\nu} - q^\mu q^\nu / q^2$, and q is the photon 4-momentum, $Q^2 = -q^2$. It is easy to see that the resulting current J'^μ obeys exactly the same WT identities as J^μ . Thus, as long as gauge-invariance is implemented at the real-photon point, the inclusion of the form factors

via Eq. (1) will give the gauge-invariant current for $Q^2 \neq 0$. For example, the bare $NN\gamma$ and $\pi\pi\gamma$ vertex functions and the Kroll-Rudermann term are:

$$\Gamma_{NN\gamma}^\mu = e\gamma^\mu + e[F_1(Q^2) - 1]O^{\mu\nu}\gamma_\nu + \frac{e\kappa_N}{2m_N}F_2(Q^2)i\sigma^{\mu\nu}q_\nu, \quad (2a)$$

$$\Gamma_{\pi\pi\gamma}^\mu = e(k+k')^\mu + e[F_\pi(Q^2) - 1]O^{\mu\nu}(k+k')_\nu, \quad (2b)$$

$$J_{KR}^\mu = \frac{eg_{\pi N}}{2m_N}\{\gamma^\mu + [F_A(Q^2) - 1]O^{\mu\nu}\gamma_\nu\}\gamma_5. \quad (2c)$$

The above procedure allows us to use the experimentally determined form factors in the Born terms. Since many of the form factors are of the dipole form we introduce: $F_D(Q^2; \Lambda^2) = (1 + Q^2/\Lambda^2)^{-2}$. The newly measured *proton electric* form factor [13], we represent by the following form: $G_E^p(Q^2) = (1 + Q^2e^{-Q^2})F_D(Q^2; 0.4)$; it is in a very good agreement with the more common form [14] in the range of Q^2 under consideration. For the *neutron electric* form factor we use the parametrization of Galster [15]. For the *magnetic* form factors of both *proton and neutron* we use the dipole form: $G_E^{p/n}(Q^2) = \mu_{p/n}F_D(Q^2, 0.71)$. For the *pion* form factor we use the monopole form: $F_\pi(Q^2) = (1 + Q^2/0.45)^{-1}$, while, for the axial form factor we use: $F_A(Q^2) = F_D(Q^2, 0.9)$. For the vector mesons (ρ/ω) we use the prediction of [16]: $F_{\rho/\omega} = (1 + Q^2)/(1 + 3.04Q^2 + 2.42Q^4 + 0.36Q^6)$.

The only undetermined form factors in our model are then the $\gamma N\Delta$ form factors. These we determine within the framework of our model by fitting to the MAID multipole analysis of data in the region of the Δ -resonance.

Let us first consider the $\gamma N\Delta$ vertex function. We write it in the Lorenz-covariant form that obeys both electromagnetic and spin-3/2 gauge symmetries (see [18, 19, 20] for details):

$$\begin{aligned} \Gamma_{\gamma\Delta N}^{\alpha\mu}(p, q) = & -\frac{3e(m_\Delta + m_N)}{2m_N[(m_\Delta + m_N)^2 - q^2]} \{g_M(Q^2)\varepsilon^{\alpha\mu\beta\nu}p_\beta q_\nu \\ & + g_E(Q^2)(p \cdot q g^{\alpha\mu} - q^\alpha p^\mu) i\gamma^5 \\ & + g_C(Q^2)(1/m_N)[q^2(p^\mu \gamma^\alpha - \not{p} g^{\alpha\mu}) + q^\mu(\not{p} q^\alpha - p \cdot q \gamma^\alpha)] i\gamma^5\}, \end{aligned} \quad (3)$$

where q (μ) and p (α) are the four-momenta (vector-indexes) of the photon and Δ respectively. This vertex function has the advantage of decoupling the unphysical spin-1/2 sector of the spin-3/2 field, which can be viewed as the result of the transversality property [19]: $p_\alpha \Gamma_{\gamma\Delta N}^{\alpha\mu} = 0$.

At the Δ -pole (the mass shell of the Δ), we can relate these couplings to the more conventional decomposition of Jones and Scadron [17] which is done in terms of G_M, G_E , and G_C where M, E , and C refer to magnetic, electric and Coulomb $\gamma N\Delta$ form factors, similar to *Sachs* form factors of the nucleon. Defining, $D(Q^2) = Q^2 + (m_\Delta - m_N)^2$ and $P(Q^2) = m_\Delta^2 - m_N^2 - Q^2$, we find the following relation between the two sets of form factors:

$$g_M = G_M - G_E \quad (4a)$$

$$g_E = \frac{2}{D}[P G_E + Q^2 G_C] \quad (4b)$$

$$g_C = \frac{m_N}{m_\Delta D}[4m_\Delta^2 G_E - P G_C] \quad (4c)$$

In the actual calculations we use the vertex function (3) with form factors expressed in terms of G_M , G_E and G_C via Eq. (4). The Δ contribution to the resonant multipoles $M_{1+}^{3/2}$, $E_{1+}^{3/2}$ and $S_{1+}^{3/2}$ are directly proportional to G_M , G_E and G_C , respectively.

To parametrize the $\gamma N\Delta$ form factors we universally use the form:

$$G_I(Q^2) = G_I \frac{1 + (Q^2/A_I) e^{-Q^2/B_I}}{(1 + Q^2/\Lambda_I^2)^2}, \quad I = M, E, C. \quad (5)$$

Here we have built in a constraint from perturbative QCD (pQCD) such that these form factors fall as Q^{-4} (modulo logs) for asymptotically large Q , see e.g. [21]. This is the principal difference with parametrizations of Refs. [9, 10] which fall off exponentially and hence do not satisfy the pQCD constraint.

The photo-couplings, i.e., G_M and G_E , are determined by the photoproduction multipoles $M_{1+}^{3/2}$ and $E_{1+}^{3/2}$ at the resonance position $W \simeq 1232$ MeV. The strength of G_C is determined by $S_{1+}^{3/2}$ at low Q^2 . We then have determined the Q^2 dependence of G_M by comparing our calculations to the experimentally extracted M_{1+} multipole at an invariant energy $W = 1232$ MeV, see Fig. 2. The solid line here represents our full model calculation of this multipole. The dotted line represents the result obtained by including only the Δ s -channel exchange in the electroproduction potential. In doing so we exclude mechanisms of resonance electroexcitation through the pion cloud, see, e.g., Fig. 3. In our model these mechanisms result from the πN rescattering through the resonant channel. Such mechanisms apparently account for about 50% of the M_{1+} strength near the photon point and about 25% near $Q^2 = 4$ GeV². This is in a qualitative agreement with the findings of Refs. [9, 10].

The parameters of the electric and Coulomb form factors are adjusted for the best description of the corresponding resonant multipoles known from MAID analysis. The values of the extracted $\gamma N\Delta$ parameters are summarized in Table I.

The ratios of the resonant multipoles: $R_{EM} = \text{Im}E_{1+}/\text{Im}M_{1+}$ and $R_{SM} = \text{Im}S_{1+}/\text{Im}M_{1+}$ carry important information about the N to Δ transition and about the admixture the D -wave component in the nucleon wave function in particular. The focus of several recent theoretical [9, 10, 23, 24] and experimental [5, 6] studies has been the extraction of the Q^2 -dependence of these ratios. This dependence can potentially tell us about the range of the momentum transfer where pQCD becomes applicable (pQCD predicts $R_{EM} = 100\%$ and $R_{SM} = \text{const}$).

In Fig. 4 we display the Q^2 -dependence of R_{EM} and R_{SM} obtained in our model and compared with a number of results of recent measurements and calculations. In particular, at the real photon point we find $R_{EM} \cong -2.7\%$ and $R_{SM} \cong -2.3\%$. As for the Q^2 dependence, we can see that R_{EM} shows a systematic tendency to cross zero in the region between 3 and 4 (GeV/c)². This is in contrast to the recent data analysis [5] or Sato and Lee model [9] which conclude that R_{EM} stays negative and is virtually flat in this domain of Q^2 . Our results are in a better agreement with the data analysis of Kamalov *et al.* [23] and Aznauryan [24], which indicate the sign change of R_{EM} below $Q^2 = 4$ (GeV/c)². Additional experiments and extractions of multipoles at higher Q^2 would be desirable to investigate this point. Our result for R_{SM} is in agreement with most of the mentioned analyses. Again the dotted curves represent the result of resonance dominance, when $R_{EM} = -G_E/G_M$ and $R_{SM} = G_C/G_M$.

After determining the Q^2 dependence of the $\gamma N\Delta$ form factors, we calculated the virtual-photon differential cross sections, spanning a wide range of Q^2 values as well as various energies W . In Fig. 5 we show several of these calculations. We find the agreement with the

recent JLab experiments quite pleasing, especially in view of the fact that the model has very few adjustable parameters, all given in Table I.

In summary, we have extended the relativistic dynamical model of Refs. [1, 2] to calculate the pion electro-production reactions. We were able to use realistic electromagnetic form factors at each photon vertex by carefully treating the problem of gauge invariance for the Born terms. For the resonance terms we parameterized the *bare* $\gamma N\Delta$ form factors such that the pQCD asymptotic Q^2 constraint is fulfilled, and then fitted them to the cross-section data and resonant multipoles of MAID up to 4 (GeV/c)². Our result for R_{EM} in this Q^2 domain is still very far from the pQCD prediction of 100%. Although, we do find that R_{EM} shows a strong tendency to cross zero and change sign in the region between 3 and 4 (GeV/c)². Our model calculation of the virtual photon cross section of $p(e, e'p)\pi^0$ shows a very good agreement with recent data up to 4 (GeV/c)² in the Δ resonance energy region. A thorough study of the existing data base of pion electroproduction reactions using this model is underway [25].

We thank Marc Vanderhaeghen for helpful discussions. This work is supported by the DOE under the contracts DE-FG02-93ER40756, DE-FG05-88ER40435, DE-FG02-93ER-40762, DE-AC05-84ER-40150, and the NSF under grant NSF-SGER-0094668.

-
- [1] V. Pascalutsa and J. A. Tjon, Phys. Rev. C **54**, 054003 (2000); Phys. Lett. B **435**, 245 (1998).
 - [2] V. Pascalutsa and J. A. Tjon, arXiv:nucl-th/0407068.
 - [3] G. Blaupied *et al.* Phys. Rev. Lett. **79**, 4337 (1997).
 - [4] R. Beck *et al.*, Phys. Rev. C **61**, 035204 (2000).
 - [5] V. V. Frolov *et al.*, Phys. Rev. Lett. **82**, 45 (1999).
 - [6] K. Joo *et al.*, Phys. Rev. Lett. **88**, 12201 (2002).
 - [7] C. Mertz *et al.*, Phys. Rev. Lett. **86**, 2963 (2001).
 - [8] D. Drechsel, O. Hanstein, S. S. Kamalov and L. Tiator, Nucl. Phys. A **645** (1999) 145.
 - [9] T. Sato and T. -S. H. Lee, Phys. Rev. C **54**, 2660 (1996); *ibid.* **63**, 055201 (2001).
 - [10] S. S. Kamalov and S. N. Yang, Phys. Rev. Lett. **83**, 4494 (1999).
 - [11] J. H. Koch, V. Pascalutsa and S. Scherer, Phys. Rev. C **65**, 045202 (2002).
 - [12] F. Gross and D. O. Riska, Phys. Rev. C **36**, 1928 (1987).
 - [13] M. K. Jones *et al.*, Phys. Rev. Lett. **84**, 1398 (2000).
 - [14] P. A. M. Guichon and M. Vanderhaeghen, Phys. Rev. Lett. **91**, 142303 (2003).
 - [15] S. Galster *et al.*, Nucl. Phys. **32B**, 221 (1971).
 - [16] P. Maris and P. C. Tandy, Phys. Rev. C **65**, 045211 (2002).
 - [17] H. F. Jones and M. D. Scadron, Annals Phys. **81**, 1 (1973).
 - [18] V. Pascalutsa, Phys. Rev. D **58**, 096002 (1998); arXiv:nucl-th/0303005.
 - [19] V. Pascalutsa and R.G.E. Timmermans, Phys. Rev. C **60**, 042201(R) (1999).
 - [20] V. Pascalutsa, Phys. Lett. B **503**, 85 (2001).
 - [21] C. E. Carlson and N. C. Mukhopadhyay, Phys. Rev. Lett. **81**, 2646 (1998).
 - [22] W. Bartel *et al.*, Phys. Lett. **28B**, 148 (1968); K. Bätzner *et al.*, *ibid.* **39B**, 575 (1972); J. C. Alder *et al.*, *ibid.* **46B**, 573 (1972); S. Stein *et al.*, Phys. Rev. D **12**, 1884 (1975); P. Stoler, *ibid.* **44**, 73 (1991); L. M. Stuart *et al.*, *ibid.* **58**, 032003 (1998).
 - [23] S. S. Kamalov *et al.*, Phys. Rev. C **64**, 032001(R) (2001).
 - [24] I. G. Aznauryan, arXiv:nucl-th/0207087.

[25] G. Caia, Ph.D. Thesis, Ohio University, 2004 (unpublished).

	$G_M(Q^2)$	$G_E(Q^2)$	$G_C(Q^2)$
$G(0)$	3.10	0.05	-0.18
Λ^2 [GeV ²]	0.6	0.5	0.8
A [GeV ²]	1.0	-1.1	-0.9
B [GeV ²]	1.2	2.0	1.0

TABLE I: Parameters of the $\gamma N \Delta$ form factors extracted in our model.




FIG. 1: The Born, vector-meson, and Δ -isobar contributions included in the electroproduction potential.




FIG. 2: Model description of $\text{Im}M_{1+}^{3/2}/F_D$, with $F_D = (1 + Q^2/0.71)^{-2}$. The data at $Q^2 = 2.8$ and 4.0 (GeV/c)^2 are taken from [5], other data are from [22]. The dotted line is the calculation including only the direct Δ exchange in $V_{\pi\gamma}$, hence leaving out resonant mechanisms due to the πN rescattering.




FIG. 3: Examples of the dynamical mechanisms of Δ -resonance electroexcitation.




FIG. 4: Results for the ratios of the resonant multipoles R_{EM} and R_{SM} . The data points are from CLAS [5, 6], LEGS [3], MAMI [4], MIT-Bates [7], and the MAID reanalysis of JLab data by Kamalov *et al.* [23]. The asterisks represent predictions of the Sato and Lee model [9].




FIG. 5: Results for the virtual-photon differential cross sections on $p(e, e'p)\pi^0$. The data points are from Refs. [5, 6].